

*Daboia russelli*.—5. Longest was 4' 6".

*Echis carinata*.—3. All these were seen bottled in Barnes' School, Deolali.

*Python molurus*.—1. This was sent alive from Peint and measured 8 feet.

*Eryx conicus*.—4.

*Typhlops brahminus*.—1. This was also seen bottled in Barnes' School, Deolali.

So out of 101 snakes 32 were highly poisonous (Cobras, Kraits, Russell's Vipers and Echis).

AHMEDNAGAR,  
August 5, 1930.

K. G. GHARPUREY,  
Lt.-Col., I.M.S.

### XXVII.—OCCURRENCE OF *CHARASIA DORSALIS* OUTSIDE THE MYSORE PLATEAU.

In part 2, Vol. xxxiii, p. 454, H. Jouguet states that '*Charasia dorsalis* does not seem to be distributed outside the Mysore Plateau'. I may state that specimens of this lizard were taken by me in a ruined temple near the Travellers' Bungalow at Gingee in the S. Arcot District of this (Madras) Presidency in the year 1920.

S. G. MANAVALA RAMANUJAN.

PRESIDENCY COLLEGE,  
MADRAS,  
August 15, 1930.

### XXVIII.—SOME MEASUREMENTS OF THE ESTUARY CROCODILE (*CROCODILUS POROSUS*) FROM SARAWAK.

In the last number of the *Journal of the Bombay Natural History Society* (Vol. XXXIV, p. 584) attention is drawn to the lack of measurements of the Estuarine Crocodile and the following will therefore be of interest :—

Sex	Total Length	Head Length	Tail Length	Total Weight	Brain Weight
1 ...	1' 4" ...	2.75" ...	3.5" ...	5 ozs.	
2 ♀ ...	1' 9" ...	3.25" ...	10" ...	10 ,,	
3 ♂ ...	1' 9" ...	3.5" ...	10.5" ...	9 ,,	
4 ♀ ...	1' 10" ...	3.4" ...	11.25" ...	11 ,,	
5 ♂ ...	1' 10.5" ...	3.5" ...	12" ...	11 ,,	18 grains.
6 ♂ ...	2' 1.5" ...	4" ...	1' 0.5" ...	1 lb.	
7 ...	2' 3" ...	4" ...	1' 3" ...	1 ,, 5 ,,	
8 ♀ ...	2' 2.5" ...	4" ...	1' 2.5" ...	1 ,, 3 ,,	